Desired Performance	Target Measures	Baseline	Ballad Health	
Ы			FY24	FYTD25
•	PSI 3 Pressure Ulcer Rate	1.07	0.36	0.34
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.05	0.08
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.05	0.00
•	PSI 8b In Hospital Fall with Other Fracture Rate		0.23	0.00
•	PSI 9 Postoperative Hemorrhage or Hematoma Rate	1.59	1.43	1.35
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76	1.58	0.00
+	PSI 11 Postoperative Respiratory Failure Rate	9.24	4.14	10.04
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31	3.06	2.06
+	PSI 13 Postoperative Sepsis Rate	3.58	3.99	5.01
+	PSI 14 Postoperative Wound Dehiscence Rate	0.83	1.29	5.71
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18	0.09	0.50
<b>1</b>	SMB: Sepsis Management Bundle	56.9%	62.9%	68.0%
+	CLABSI	0.711	0.782	0.799
•	CAUTI	0.558	0.821	0.809
•	SSI COLON Surgical Site Infection	2.13	3.70	1.52
•	SSI HYST Surgical Site Infection	0.71	3.37	0.00
•	MRSA	0.047	0.051	0.071
•	CDIFF	0.671	0.110	0.179
<b>1</b>	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	76.8	77.9
<b>1</b>	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	77.2	77.7
•	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	59.6	57.7
1	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	85.9	84.5
<b>1</b>	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	47.6	47.6
•	READM30HF Heart Failure 30day readmissions rate	27.7%	24.1%	23.6%
+	READM30PN Pneumonia 30day readmission rate	17.9%	18.0%	25.2%
<b>+</b>	Sepsis In House Mortality	10.7%	8.5%	8.5%
•	MORT30HF Heart failure 30day mortality rate	3.0%	3.2%	2.2%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	4.5%	4.7%
•	Left without being seen	0.83%	0.83%	0.73%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	151.8	154.0
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	432.0	376.5

Desired Performance	Target Measures	Baseline	Johnson City Med Center	
ď			FY24	FYTD25
•	PSI 3 Pressure Ulcer Rate	1.07	0.55	1.06
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.14	0.28
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.05	0.00
+	PSI 8b In Hospital Fall with Other Fracture Rate		0.23	0.00
•	PSI 9 Postoperative Hemorrhage or Hematoma Rate	1.59	1.01	0.00
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76	1.16	0.00
•	PSI 11 Postoperative Respiratory Failure Rate	9.24	3.15	11.95
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31	2.69	2.22
+	PSI 13 Postoperative Sepsis Rate	3.58	4.82	0.00
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83	3.61	7.69
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18	0.00	0.00
<b>1</b>	SMB: Sepsis Management Bundle	56.9%	40.5%	41.7%
•	CLABSI	0.711	0.618	1.153
•	CAUTI	0.558	1.357	1.672
•	SSI COLON Surgical Site Infection	2.13	7.87	20.00
•	SSI HYST Surgical Site Infection	0.71	2.63	0.00
•	MRSA	0.047	0.065	0.125
•	CDIFF	0.671	0.089	0.139
<b>1</b>	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	71.4	73.2
<b>1</b>	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	71.4	73.0
•	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	52.6	46.9
<b>↑</b>	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	82.6	80.7
<b>1</b>	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	40.4	40.5
•	READM30HF Heart Failure 30day readmissions rate	27.7%	27.0%	40.5%
•	READM30PN Pneumonia 30day readmission rate	17.9%	17.5%	26.9%
•	Sepsis In House Mortality	10.7%	13.8%	17.4%
•	MORT30HF Heart failure 30day mortality rate	3.0%	3.7%	2.3%
<b>+</b>	MORT30PN Pneumonia 30day mortality rate	5.0%	6.1%	10.8%
+	Left without being seen	0.83%	0.76%	0.62%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	209.0	192.5
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	617.5	839.0

Desired Performance	Target Measures	Baseline		n Valley I Center
D Perf			FY24	FYTD25
•	PSI 3 Pressure Ulcer Rate	1.07	0.20	0.00
+	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00
•	PSI 8b In Hospital Fall with Other Fracture Rate		0.14	0.00
•	PSI 9 Postoperative Hemorrhage or Hematoma Rate	1.59	0.71	4.30
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76	4.12	0.00
•	PSI 11 Postoperative Respiratory Failure Rate	9.24	6.44	8.16
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31	3.34	0.00
•	PSI 13 Postoperative Sepsis Rate	3.58	4.29	8.20
<b>+</b>	PSI 14 Postoperative Wound Dehiscence Rate	0.83	0.00	15.63
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18	0.50	0.00
<b>1</b>	SMB: Sepsis Management Bundle	56.9%	42.3%	50.0%
•	CLABSI	0.711	1.296	0.603
•	CAUTI	0.558	1.294	0.000
•	SSI COLON Surgical Site Infection	2.13	5.94	0.00
<b>+</b>	SSI HYST Surgical Site Infection	0.71	8.00	0.00
•	MRSA	0.047	0.087	0.079
•	CDIFF	0.671	0.186	0.237
•	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	74.7	74.3
•	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	77.5	77.2
<b>1</b>	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	58.5	57.0
1	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	86.8	82.0
<b>1</b>	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	46.5	43.8
•	READM30HF Heart Failure 30day readmissions rate	27.7%	22.7%	19.4%
+	READM30PN Pneumonia 30day readmission rate	17.9%	17.9%	29.0%
•	Sepsis In House Mortality	10.7%	11.9%	9.7%
•	MORT30HF Heart failure 30day mortality rate	3.0%	3.9%	4.5%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	5.7%	4.1%
•	Left without being seen	0.83%	0.42%	0.64%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	216.8	217.5
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	553.1	438.5

Desired Performance	Target Measures	Bristol Re Baseline Medical (		_
Pe			FY24	FYTD25
+	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.00
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.15	0.00
•	PSI 8b In Hospital Fall with Other Fracture Rate		0.30	0.00
•	PSI 9 Postoperative Hemorrhage or Hematoma Rate	1.59	0.49	0.00
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76	0.00	0.00
•	PSI 11 Postoperative Respiratory Failure Rate	9.24	7.40	0.00
+	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31	2.74	2.98
•	PSI 13 Postoperative Sepsis Rate	3.58	3.07	0.00
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83	0.00	0.00
+	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18	0.00	0.00
<b>1</b>	SMB: Sepsis Management Bundle	56.9%	61.0%	66.7%
+	CLABSI	0.711	0.240	0.690
•	CAUTI	0.558	0.712	0.641
•	SSI COLON Surgical Site Infection	2.13	2.65	0.00
•	SSI HYST Surgical Site Infection	0.71	2.70	0.00
•	MRSA	0.047	0.042	0.079
•	CDIFF	0.671	0.152	0.158
<b>1</b>	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	76.8	77.0
1	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	76.2	75.9
<b>1</b>	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	62.2	61.7
<b>1</b>	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	87.5	84.5
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	48.8	47.5
•	READM30HF Heart Failure 30day readmissions rate	27.7%	25.4%	16.1%
•	READM30PN Pneumonia 30day readmission rate	17.9%	19.5%	20.0%
•	Sepsis In House Mortality	10.7%	8.8%	7.1%
•	MORT30HF Heart failure 30day mortality rate	3.0%	2.5%	1.7%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	6.0%	5.5%
•	Left without being seen	0.83%	1.08%	1.11%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	193.0	209.0
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	379.3	349.0

Desired Performance	Target Measures	Baseline		Memorial pital
ď			FY24	FYTD25
<b>+</b>	PSI 3 Pressure Ulcer Rate	1.07	1.10	0.00
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.18	0.00
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00
•	PSI 8b In Hospital Fall with Other Fracture Rate		0.18	0.00
•	PSI 9 Postoperative Hemorrhage or Hematoma Rate	1.59	3.60	0.00
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76	0.00	0.00
•	PSI 11 Postoperative Respiratory Failure Rate	9.24	0.00	0.00
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31	4.51	0.00
•	PSI 13 Postoperative Sepsis Rate	3.58	0.00	0.00
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83	0.00	0.00
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18	0.00	0.00
•	SMB: Sepsis Management Bundle	56.9%	66.4%	72.7%
•	CLABSI	0.711	1.730	3.021
•	CAUTI	0.558	0.235	1.616
•	SSI COLON Surgical Site Infection	2.13	1.11	0.00
•	SSI HYST Surgical Site Infection	0.71	0.00	
•	MRSA	0.047	0.034	0.000
•	CDIFF	0.671	0.035	0.000
•	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	75.2	80.0
<b>1</b>	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	79.0	82.6
<b>1</b>	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	57.7	62.5
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	86.6	88.3
<b>1</b>	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	47.6	48.7
•	READM30HF Heart Failure 30day readmissions rate	27.7%	22.6%	25.0%
•	READM30PN Pneumonia 30day readmission rate	17.9%	18.0%	30.2%
•	Sepsis In House Mortality	10.7%	10.1%	7.9%
•	MORT30HF Heart failure 30day mortality rate	3.0%	3.3%	2.3%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	5.0%	3.0%
•	Left without being seen	0.83%	1.65%	0.78%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	212.8	158.5
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	409.5	378.0

Desired Performance	Target Measures	Baseline		woods ty Hospital
Per			FY24	FYTD25
	PSI 3 Pressure Ulcer Rate	1.07	0.32	0.00
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.18	0.00
•	PSI 8b In Hospital Fall with Other Fracture Rate		0.54	0.00
	PSI 9 Postoperative Hemorrhage or Hematoma Rate	1.59	3.23	5.68
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76	0.00	0.00
•	PSI 11 Postoperative Respiratory Failure Rate	9.24	0.00	33.33
+	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31	3.03	5.03
•	PSI 13 Postoperative Sepsis Rate	3.58	2.15	24.39
+	PSI 14 Postoperative Wound Dehiscence Rate	0.83	0.00	11.76
	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18	0.00	3.44
<b>1</b>	SMB: Sepsis Management Bundle	56.9%	65.1%	62.5%
•	CLABSI	0.711	0.968	0.000
•	CAUTI	0.558	0.448	0.000
•	SSI COLON Surgical Site Infection	2.13	2.03	0.00
•	SSI HYST Surgical Site Infection	0.71	0.00	0.00
•	MRSA	0.047	0.036	0.000
•	CDIFF	0.671	0.081	0.000
<b>†</b>	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	82.9	83.0
•	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	80.6	80.7
•	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	64.7	59.5
1	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	88.8	87.3
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	56.4	57.6
•	READM30HF Heart Failure 30day readmissions rate	27.7%	18.9%	0.0%
•	READM30PN Pneumonia 30day readmission rate	17.9%	16.2%	21.4%
•	Sepsis In House Mortality	10.7%	2.5%	4.2%
•	MORT30HF Heart failure 30day mortality rate	3.0%	3.6%	0.0%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	2.7%	2.8%
•	Left without being seen	0.83%	1.06%	1.19%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	187.8	204.0
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	624.8	485.5

Desired Performance	Target Measures	Baseline		n Path ty Hospital
Pe			FY24	FYTD25
•	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.00
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00
•	PSI 8b In Hospital Fall with Other Fracture Rate		0.00	0.00
•	PSI 9 Postoperative Hemorrhage or Hematoma Rate	1.59	0.00	0.00
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76	0.00	0.00
•	PSI 11 Postoperative Respiratory Failure Rate	9.24	0.00	0.00
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31	4.88	0.00
•	PSI 13 Postoperative Sepsis Rate	3.58	0.00	0.00
+	PSI 14 Postoperative Wound Dehiscence Rate	0.83	9.35	0.00
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18	0.00	0.00
<b>1</b>	SMB: Sepsis Management Bundle	56.9%	75.6%	83.3%
•	CLABSI	0.711	0.000	0.000
•	CAUTI	0.558	0.000	0.000
•	SSI COLON Surgical Site Infection	2.13	0.00	0.00
•	SSI HYST Surgical Site Infection	0.71	0.00	0.00
•	MRSA	0.047	0.000	0.000
•	CDIFF	0.671	0.000	0.000
•	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	79.3	93.6
•	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	81.1	86.2
•	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	61.1	56.3
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	84.8	90.9
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	50.1	58.4
<b>+</b>	READM30HF Heart Failure 30day readmissions rate	27.7%	12.5%	0.0%
<b>+</b>	READM30PN Pneumonia 30day readmission rate	17.9%	16.3%	26.3%
<b>+</b>	Sepsis In House Mortality	10.7%	0.5%	0.0%
•	MORT30HF Heart failure 30day mortality rate	3.0%	0.0%	0.0%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	0.0%	0.0%
<b>+</b>	Left without being seen	0.83%	0.34%	0.50%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	152.8	170.0
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	433.0	650.0

Desired Performance	Target Measures	Baseline		ommunity pital
Pe			FY24	FYTD25
•	PSI 3 Pressure Ulcer Rate	1.07	0.62	0.00
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00
•	PSI 8b In Hospital Fall with Other Fracture Rate		0.00	0.00
•	PSI 9 Postoperative Hemorrhage or Hematoma Rate	1.59	0.00	0.00
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76	0.00	0.00
•	PSI 11 Postoperative Respiratory Failure Rate	9.24	0.00	0.00
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31	3.01	0.00
•	PSI 13 Postoperative Sepsis Rate	3.58	0.00	0.00
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83	0.00	0.00
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18	0.00	0.00
•	SMB: Sepsis Management Bundle	56.9%	64.2%	25.0%
•	CLABSI	0.711	0.000	0.000
•	CAUTI	0.558	0.000	0.000
•	SSI COLON Surgical Site Infection	2.13	2.86	0.00
+	SSI HYST Surgical Site Infection	0.71	0.00	
+	MRSA	0.047	0.000	0.000
•	CDIFF	0.671	0.087	0.000
<b>1</b>	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	77.6	76.6
<b>1</b>	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	78.0	72.8
•	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	59.9	60.3
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	82.8	82.5
<b>1</b>	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	49.0	52.4
•	READM30HF Heart Failure 30day readmissions rate	27.7%	25.2%	18.2%
•	READM30PN Pneumonia 30day readmission rate	17.9%	16.8%	20.0%
+	Sepsis In House Mortality	10.7%	4.1%	7.2%
•	MORT30HF Heart failure 30day mortality rate	3.0%	2.4%	0.0%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	3.1%	8.1%
•	Left without being seen	0.83%	0.86%	0.51%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	150.7	159.5
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	612.3	375.0

Desired Performance	Target Measures	Baseline		neville ty Hospital
Per			FY24	FYTD25
•	PSI 3 Pressure Ulcer Rate	1.07	0.72	0.00
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00
•	PSI 8b In Hospital Fall with Other Fracture Rate		0.45	0.00
•	PSI 9 Postoperative Hemorrhage or Hematoma Rate	1.59	5.15	0.00
+	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76	0.00	0.00
•	PSI 11 Postoperative Respiratory Failure Rate	9.24	5.78	0.00
+	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31	4.84	7.19
•	PSI 13 Postoperative Sepsis Rate	3.58	6.71	0.00
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83	0.00	0.00
	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18	0.00	0.00
•	SMB: Sepsis Management Bundle	56.9%	58.1%	80.0%
•	CLABSI	0.711	1.021	0.000
•	CAUTI	0.558	0.000	0.000
•	SSI COLON Surgical Site Infection	2.13	4.38	0.00
•	SSI HYST Surgical Site Infection	0.71	0.00	
•	MRSA	0.047	0.000	0.000
•	CDIFF	0.671	0.048	0.270
•	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	78.9	83.9
•	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	77.1	84.9
•	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	63.2	67.1
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	87.7	87.0
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	45.9	51.6
•	READM30HF Heart Failure 30day readmissions rate	27.7%	25.4%	15.4%
•	READM30PN Pneumonia 30day readmission rate	17.9%	18.9%	20.0%
•	Sepsis In House Mortality	10.7%	5.8%	4.5%
•	MORT30HF Heart failure 30day mortality rate	3.0%	3.4%	2.0%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	3.0%	0.0%
•	Left without being seen	0.83%	0.92%	0.71%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	163.0	212.0
	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	633.9	576.5

Desired Performance	Target Measures	Baseline		County ty Hospital
De	raiget measures	Baseinie	Communi	
			FY24	FYTD25
•	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.00
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00
•	PSI 8b In Hospital Fall with Other Fracture Rate		0.87	0.00
•	PSI 9 Postoperative Hemorrhage or Hematoma Rate	1.59	0.00	0.00
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76	0.00	0.00
•	PSI 11 Postoperative Respiratory Failure Rate	9.24	0.00	0.00
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31	0.00	0.00
•	PSI 13 Postoperative Sepsis Rate	3.58	0.00	0.00
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83	0.00	0.00
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration	1.18	0.00	0.00
•	SMB: Sepsis Management Bundle	56.9%	86.2%	66.7%
-	CLABSI	0.711	0.000	0.000
•	CAUTI	0.558	0.000	0.000
•	SSI COLON Surgical Site Infection	2.13	0.000	0.000
•	SSI HYST Surgical Site Infection	0.71		
•	MRSA	0.047	0.000	0.000
•	CDIFF	0.671	0.000	0.000
<b>1</b>	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	78.2	75.9
<b>1</b>	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	79.7	81.0
<b>1</b>	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	60.4	54.2
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	84.0	95.8
<b>1</b>	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	47.9	40.1
•	READM30HF Heart Failure 30day readmissions rate	27.7%	15.4%	20.0%
•	READM30PN Pneumonia 30day readmission rate	17.9%	16.6%	20.0%
•	Sepsis In House Mortality	10.7%	4.6%	0.0%
•	MORT30HF Heart failure 30day mortality rate	3.0%	5.5%	0.0%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	2.3%	0.0%
•	Left without being seen	0.83%	1.02%	0.58%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	144.0	121.5
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	658.0	301.5

Desired Performance	Target Measures	Baseline		re Shoals pital
A.			FY24	FYTD25
<b>+</b>	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.00
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00
•	PSI 8b In Hospital Fall with Other Fracture Rate		0.00	0.00
<b>+</b>	PSI 9 Postoperative Hemorrhage or Hematoma Rate	1.59	2.78	0.00
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76	0.00	0.00
<b>+</b>	PSI 11 Postoperative Respiratory Failure Rate	9.24	0.00	0.00
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31	0.00	0.00
•	PSI 13 Postoperative Sepsis Rate	3.58	7.35	0.00
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83	0.00	0.00
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18	0.00	0.00
<b>1</b>	SMB: Sepsis Management Bundle	56.9%	60.8%	66.7%
•	CLABSI	0.711	0.000	0.000
•	CAUTI	0.558	0.000	6.098
•	SSI COLON Surgical Site Infection	2.13	6.67	0.00
•	SSI HYST Surgical Site Infection	0.71	15.38	0.00
•	MRSA	0.047	0.000	0.000
•	CDIFF	0.671	0.000	0.986
<b>1</b>	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	76.4	82.4
<b>1</b>	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	79.5	85.2
<b>1</b>	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	58.7	70.3
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	80.8	91.9
<b>1</b>	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	43.9	52.8
•	READM30HF Heart Failure 30day readmissions rate	27.7%	17.5%	16.7%
<b>+</b>	READM30PN Pneumonia 30day readmission rate	17.9%	21.0%	30.8%
<b>+</b>	Sepsis In House Mortality	10.7%	3.0%	6.8%
•	MORT30HF Heart failure 30day mortality rate	3.0%	1.9%	0.0%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	2.5%	6.1%
•	Left without being seen	0.83%	1.25%	0.73%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	173.2	157.0
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	621.5	1177.0

Desired Performance	Target Measures	Baseline		me Pine spital
Ğ			FY24	FYTD25
•	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.00
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00
•	PSI 8b In Hospital Fall with Other Fracture Rate		0.00	0.00
•	PSI 9 Postoperative Hemorrhage or Hematoma Rate	1.59	0.00	
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76		
•	PSI 11 Postoperative Respiratory Failure Rate	9.24		
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31	0.00	
•	PSI 13 Postoperative Sepsis Rate	3.58		
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83	0.00	
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration	1.18	0.00	
<b>1</b>	SMB: Sepsis Management Bundle	56.9%	63.3%	
	CLABSI	0.711	0.000	0.000
	CAUTI	0.558	0.000	0.000
•	SSI COLON Surgical Site Infection	2.13		
•	SSI HYST Surgical Site Infection	0.71		
	MRSA	0.047	0.000	0.000
<b>+</b>	CDIFF	0.671	0.000	0.000
<b>1</b>	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	82.4	83.3
<b>1</b>	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	79.9	50.0
•	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	61.5	50.0
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	91.0	50.0
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	47.7	50.0
•	READM30HF Heart Failure 30day readmissions rate	27.7%	12.0%	33.3%
<b>+</b>	READM30PN Pneumonia 30day readmission rate	17.9%	17.6%	33.3%
•	Sepsis In House Mortality	10.7%	1.9%	0.0%
•	MORT30HF Heart failure 30day mortality rate	3.0%	0.0%	0.0%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	1.9%	0.0%
•	Left without being seen	0.83%	0.61%	0.43%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	134.6	151.0
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	322.1	285.0

Desired Performance	Target Measures	Baseline	Hawkins County Memorial Hospital	
Pe			FY24	FYTD25
•	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.00
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00
•	PSI 8b In Hospital Fall with Other Fracture Rate		0.00	0.00
•	PSI 9 Postoperative Hemorrhage or Hematoma Rate	1.59		
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76		
•	PSI 11 Postoperative Respiratory Failure Rate	9.24		
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31		
•	PSI 13 Postoperative Sepsis Rate	3.58		
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83		
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration	1.18		
•	SMB: Sepsis Management Bundle	56.9%	75.7%	100.0%
•	CLABSI	0.711	0.000	0.000
•	CAUTI	0.558	0.000	0.000
•	SSI COLON Surgical Site Infection	2.13		
•	SSI HYST Surgical Site Infection	0.71		
•	MRSA	0.047	0.000	0.000
•	CDIFF	0.671	0.000	4.975
•	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	88.3	86.7
<b>1</b>	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	85.6	53.3
<b>1</b>	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	80.8	66.7
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	86.5	63.3
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	52.4	39.1
•	READM30HF Heart Failure 30day readmissions rate	27.7%	29.6%	0.0%
<b>♣</b>	READM30PN Pneumonia 30day readmission rate	17.9%	5.5%	0.0%
•	Sepsis In House Mortality	10.7%	5.4%	12.5%
•	MORT30HF Heart failure 30day mortality rate	3.0%	0.0%	0.0%
+	MORT30PN Pneumonia 30day mortality rate	5.0%	1.8%	0.0%
•	Left without being seen	0.83%	0.21%	0.23%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	105.6	84.0
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	240.1	287.0

Desired Performance	Target Measures	Baseline	Russell County Hospital	
Ä			FY24	FYTD25
•	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.00
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00
+	PSI 8b In Hospital Fall with Other Fracture Rate		0.00	0.00
•	PSI 9 Postoperative Hemorrhage or Hematoma Rate	1.59		
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76		
•	PSI 11 Postoperative Respiratory Failure Rate	9.24		
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31		
•	PSI 13 Postoperative Sepsis Rate	3.58		
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83		
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18	0.00	0.00
•	SMB: Sepsis Management Bundle	56.9%	76.9%	88.9%
+	CLABSI	0.711	0.000	0.000
•	CAUTI	0.558	0.000	0.000
•	SSI COLON Surgical Site Infection	2.13		
•	SSI HYST Surgical Site Infection	0.71		
•	MRSA	0.047	0.000	0.000
+	CDIFF	0.671	0.000	0.000
•	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	84.2	81.3
<b>1</b>	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	79.9	70.3
<b>1</b>	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	62.9	50.0
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	87.7	72.1
<b>1</b>	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	44.9	54.1
•	READM30HF Heart Failure 30day readmissions rate	27.7%	34.0%	0.0%
<b>+</b>	READM30PN Pneumonia 30day readmission rate	17.9%	23.6%	25.0%
<b>+</b>	Sepsis In House Mortality	10.7%	0.5%	2.8%
•	MORT30HF Heart failure 30day mortality rate	3.0%	2.1%	0.0%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	0.7%	0.0%
•	Left without being seen	0.83%	1.14%	0.67%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	124.1	113.0
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	213.2	243.5

Desired Performance	Target Measures	Baseline	Unicoi County Hospital	
Ä			FY24	FYTD25
•	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.00
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00
+	PSI 8b In Hospital Fall with Other Fracture Rate		0.00	0.00
•	PSI 9 Postoperative Hemorrhage or Hematoma Rate	1.59		
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76		
•	PSI 11 Postoperative Respiratory Failure Rate	9.24		
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31		
+	PSI 13 Postoperative Sepsis Rate	3.58		
+	PSI 14 Postoperative Wound Dehiscence Rate	0.83		
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration	1.18	0.00	
•	SMB: Sepsis Management Bundle	56.9%	71.7%	100.0%
+	CLABSI	0.711	0.000	0.000
+	CAUTI	0.558	0.000	0.000
+	SSI COLON Surgical Site Infection	2.13		
•	SSI HYST Surgical Site Infection	0.71		
•	MRSA	0.047	0.000	0.000
•	CDIFF	0.671	0.487	0.000
<b>1</b>	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	83.7	80.7
<b>1</b>	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	81.5	78.2
<b>1</b>	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	68.4	56.4
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	89.7	92.9
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	53.8	55.9
•	READM30HF Heart Failure 30day readmissions rate	27.7%	17.9%	50.0%
•	READM30PN Pneumonia 30day readmission rate	17.9%	20.2%	33.3%
<b>+</b>	Sepsis In House Mortality	10.7%	1.5%	0.0%
•	MORT30HF Heart failure 30day mortality rate	3.0%	0.0%	0.0%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	2.2%	0.0%
•	Left without being seen	0.83%	0.41%	0.72%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	137.9	134.5
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	648.7	582.0

Desired Performance	Target Measures	Baseline Cor		Lee County Community Hospital	
D Perf			FY24	FYTD25	
•	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.00	
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00	
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00	
•	PSI 8b In Hospital Fall with Other Fracture Rate	0.00	0.00	0.00	
•	PSI 9 Postoperative Hemorrhage or Hematoma Rate	1.59	0.00	0.00	
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76			
+	PSI 11 Postoperative Respiratory Failure Rate	9.24			
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31			
•	PSI 13 Postoperative Sepsis Rate	3.58			
•	PSI 14 Postoperative Sepsis Nate	0.83			
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18			
•	SMB: Sepsis Management Bundle	56.9%	71.0%	100.0%	
-	CLABSI	0.711			
•	CAUTI	0.558			
•	SSI COLON Surgical Site Infection	2.13			
•	SSI HYST Surgical Site Infection	0.71			
•	MRSA	0.047			
•	CDIFF	0.671			
<b>1</b>	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	89.7	100.0	
<b>1</b>	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	85.3	100.0	
<b>1</b>	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	71.7	91.7	
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	80.3	95.0	
<b>1</b>	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	57.2	75.0	
•	READM30HF Heart Failure 30day readmissions rate	27.7%	12.5%		
•	READM30PN Pneumonia 30day readmission rate	17.9%	16.0%	0.0%	
•	Sepsis In House Mortality	10.7%	3.2%	0.0%	
•	MORT30HF Heart failure 30day mortality rate	3.0%	0.0%	0.0%	
•	MORT30PN Pneumonia 30day mortality rate	5.0%	0.0%	0.0%	
•	Left without being seen	0.83%	0.29%	0.68%	
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	109.0	111.5	
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	318.1	245.5	

Desired Performance	Target Measures	Baseline Con		Dickenson Community Hospital	
			FY24	FYTD25	
•	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.00	
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00	
+	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00	
+	PSI 8b In Hospital Fall with Other Fracture Rate	0.00	0.00	0.00	
•	PSI 9 Postoperative Hemorrhage or Hematoma Rate	1.59	0.00	0.00	
<b>+</b>	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76			
•	PSI 11 Postoperative Respiratory Failure Rate	9.24			
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31			
•	PSI 13 Postoperative Sepsis Rate	3.58			
+	PSI 14 Postoperative Wound Dehiscence Rate	0.83			
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18			
•	SMB: Sepsis Management Bundle	56.9%			
<u>-</u>	CLABSI	0.711			
•	CAUTI	0.558			
•	SSI COLON Surgical Site Infection	2.13			
•	SSI HYST Surgical Site Infection	0.71			
•	MRSA	0.047			
<u> </u>	CDIFF	0.671			
<b>1</b>	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	72.2		
•	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	38.9		
<b>1</b>	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	50.0		
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	75.0		
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	40.0		
•	READM30HF Heart Failure 30day readmissions rate	27.7%			
•	READM30PN Pneumonia 30day readmission rate	17.9%	0.0%		
•	Sepsis In House Mortality	10.7%			
•	MORT30HF Heart failure 30day mortality rate	3.0%			
<b>+</b>	MORT30PN Pneumonia 30day mortality rate	5.0%	0.0%		
•	Left without being seen	0.83%	0.43%	0.54%	
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	110.5	106.5	
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	161.0		

Desired Performance	Target Measures			ock County ospital	
A.			FY24	FYTD25	
•	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.00	
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00	
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00	
+	PSI 8b In Hospital Fall with Other Fracture Rate		0.00	0.00	
•	PSI 9 Postoperative Hemorrhage or Hematoma Rate	1.59			
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76			
•	PSI 11 Postoperative Respiratory Failure Rate	9.24			
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31			
•	PSI 13 Postoperative Sepsis Rate	3.58			
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83			
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18			
•	SMB: Sepsis Management Bundle	56.9%	85.7%	100.0%	
•	CLABSI	0.711			
•	CAUTI	0.558			
•	SSI COLON Surgical Site Infection	2.13			
•	SSI HYST Surgical Site Infection	0.71			
•	MRSA	0.047			
<b>+</b>	CDIFF	0.671			
•	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	84.1	91.7	
•	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	83.3	91.7	
•	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	42.5	83.3	
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	92.9	100.0	
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	70.9	61.1	
•	READM30HF Heart Failure 30day readmissions rate	27.7%	0.0%		
•	READM30PN Pneumonia 30day readmission rate	17.9%	0.0%	0.0%	
•	Sepsis In House Mortality	10.7%	5.0%	0.0%	
•	MORT30HF Heart failure 30day mortality rate	3.0%	0.0%		
+	MORT30PN Pneumonia 30day mortality rate	5.0%	9.1%	0.0%	
•	Left without being seen	0.83%	0.14%	0.00%	
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	126.9	145.0	
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	212.0	199.5	

Desired Performance	Target Measures	Baseline	Johnson Count	•
Pe			FY24	FYTD25
•	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.00
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00
+	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00
+	PSI 8b In Hospital Fall with Other Fracture Rate		0.00	0.00
•	PSI 9 Postoperative Hemorrhage or Hematoma Rate	1.59		
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76		
•	PSI 11 Postoperative Respiratory Failure Rate	9.24		
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31		
•	PSI 13 Postoperative Sepsis Rate	3.58		
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83		
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration	1.18		
<b>1</b>	SMB: Sepsis Management Bundle	56.9%		
•	CLABSI	0.711		
+	CAUTI	0.558		
+	SSI COLON Surgical Site Infection	2.13		
•	SSI HYST Surgical Site Infection	0.71		
+	MRSA	0.047		
•	CDIFF	0.671		
<b>1</b>	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	88.9	
<b>1</b>	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	66.7	
•	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	40.0	
<b>1</b>	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	45.0	
<b>1</b>	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	27.2	
•	READM30HF Heart Failure 30day readmissions rate	27.7%	0.0%	
•	READM30PN Pneumonia 30day readmission rate	17.9%	16.7%	
•	Sepsis In House Mortality	10.7%	0.0%	0.0%
•	MORT30HF Heart failure 30day mortality rate	3.0%	0.0%	
•	MORT30PN Pneumonia 30day mortality rate	5.0%	0.0%	
<b>+</b>	Left without being seen	0.83%	0.86%	1.36%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	136.7	113.0
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	770.0	